

Author declaration

Acknowledgements

Contributions from Prof. Vajira Weerasinghe, Department of Physiology, Faculty of Medicine, University of Peradeniya, Sri Lanka; Dr. Saman Senanayake, Consultant Eye Surgeon, National Hospital, Kandy, Sri Lanka; Prof. Mike Sedgwick, Consultant Neurophysiologist, Eastleigh, United Kingdom; Prof. Tharaka Dassanayake, Department of Physiology, Faculty of Medicine, University of Peradeniya, Sri Lanka; Prof. Indika Gawarammana, Department of Medicine, Faculty of Medicine, University of Peradeniya, Sri Lanka; Prof. Manoji Pathirage, Department of Medicine, Faculty of Medicine, University of Peradeniya, Sri Lanka and Prof. Graham Holder, Consultant in Clinical Electrophysiology, Moorfields Eye Hospital, United Kingdom are greatly acknowledged.

References

1. Bourne R., Steinmetz J., Flaxman S. et al. Trends in prevalence of blindness and distance and near vision impairment over 30 years: an analysis for the Global Burden of Disease Study. *Lancet Glob Heal*. 2020. doi: [https://doi.org/10.1016/S2214-109X\(20\)30425-3](https://doi.org/10.1016/S2214-109X(20)30425-3).
2. Gilbert C, Edusuriya K, Murthy GVS, Pant HB, Schmidt E, Senanayake S, et al. Prevalence, causes and magnitude of functional low vision in Sri Lanka: results from a national population based survey. *Ceylon Medical Journal*. 2018;63(5):s40–s4. . doi: DOI: <http://doi.org/10.4038/cmj.v63i5.8741>.
3. Dahanayake P, Dassanayake TL, Pathirage M, Senanayake S, Sedgwick M, Weerasinghe V. Electro-oculography in bilateral optic neuropathy. *BMC Research Notes*. 2020;13(1):287. doi: 10.1186/s13104-020-05131-0.
4. Sedgwick M, Dahanayake D, Dassanayake T, Senanayake S, Weerasinghe V. The Corneo-Fundal Potential in Bilateral Optic Neuritis: British Society for Clinical Neurophysiology (BSCN); 2019. Available from: <https://www.bscn.org.uk/abstract-details.aspx?id=113>.
5. Dahanayake D, Pathirage L, Hangilipola A, Baminiwattha D, Senanayake S, VS W. Combined use of pattern electroretinography (PERG) and pattern reversal visual evoked potentials (PVEP) in optic neuropathies. *Journal of the college of ophthalmologists of Sri Lanka*. 2017;23(2):55-8.
6. Matsumoto H, Shibasaki K, Uchigashima M, Koizumi A, Kurachi M, Moriwaki Y, et al. Localization of acetylcholine-related molecules in the retina: implication of the communication from photoreceptor to retinal pigment epithelium. *PloS one*. 2012;7(8):e42841-e. Epub 08/03. doi: 10.1371/journal.pone.0042841. PubMed PMID: 22880119.
7. Dahanayake P, Dassanayake TL, Pathirage M, Colombage A, Gawarammana IB, Senanayake S, et al. Dysfunction in macula, retinal pigment epithelium and post retinal pathway in acute organophosphorus poisoning. *Clinical Toxicology*. 2020:1-7. doi: 10.1080/15563650.2020.1771359.
8. Dahanayake P, Dassanayake TL, Pathirage M, Senanayake S, Sedgwick M, Weerasinghe V. Detection of early diabetic retinopathy using visual electrophysiological tests. *International Journal of Diabetes in Developing Countries*. 2021. doi: 10.1007/s13410-020-00916-2.
9. Dahanayake DWP., Pathirage LPMMK., Shiroma RGL., Senanayake S., VS W, editors. Combined use of pattern electroretinography (PERG) and electro-oculography (EOG) in retinal pigment epithelial disorders: preliminary data. Peradeniya university international research sessions (iPurse); 2017.
10. Dahanayake DWP., Pathirage LPMMK., Shiroma RGL., Senanayake S., VS W, editors. A study of electro-oculography (EOG) in patients with retinal disorders. 39th Annual academic sessions of the Kandy society of Medicine; 2017.